

## Information sheet for customer

# Tsumura-Kampo Choreito[猪苓湯] Extract Granules A

【Category 2 OTC medicines】

## INDICATIONS

Following symptoms in persons who have abnormality of urination, and sometimes experience thirst. This medicine can be applied to everyone regardless of physical performance:

Urination difficulty, urination pain, feeling of residual urine, frequent urination, and edema.



## ! CAUTION FOR USE

### ●Who should seek consultation

- The following persons should consult a physician, pharmacist, or registered distributor before taking this medicine.
  - Persons undergoing medical treatment from a physician.
  - Pregnant women or women suspected of being pregnant.
- The following symptoms may be adverse reactions to this medication. If any of these symptoms occur, stop taking this medicine immediately and consult a physician, pharmacist, or registered distributor. Take the instruction leaflet with you.

Related areas	Symptoms
Skin	Rash, redness, itching

- Stop taking this medicine and consult a physician, pharmacist or registered distributor if symptoms do not improve after taking this medicine for about 1 month. Take this leaflet with you.

## DOSAGE

Take the following dosages with water or warm water before a meal.

Age	One dosage	Daily dosage
15 years and over	1 packet (1.875 g)	2 times a day
7-14 years	2/3 packet	
4-6 years	1/2 packet	
2-3 years	1/3 packet	
Less than 2 years	Do not use	

[Cautions] This medication should be given to children only under adult supervision

## INGREDIENTS

2 packets (3.75 g) of this medicine contain 1.25 g of Choreito extract of the following mixed crude drugs;

JP Alminum Silicate Hydrate with Silicon Dioxide	1.5 g
JP Alisma Rhizome	1.5 g
JP Polyporus Sclerotium	1.5 g
JP Poria Sclerotium	1.5 g
Donkey Glue	1.5 g

(JP: The Japanese Pharmacopoeia)

Inactive ingredients: JP Magnesium Stearate, JP Lactose Hydrate.

## PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Store in a cool, dry place free from direct sunlight.
- (2) Keep all medication out of the reach of children.
- (3) When you take this medicine in divided dose, fold over the opened edge of the packet and store it.  
The remaining medicine should be taken within 2 days.
- (4) Since the medicine contains crude drugs, the granules color may vary for individual products, however this has no influence on the efficacy of this medicine.
- (5) Do not take this medicine after the expiration date.

*Production of this product has been approved for domestic sales in accordance with  
the Pharmaceutical Affairs Law of Japan.*



**Tsumura&Co.** 2-17-11, Akasaka, Minato-ku, Tokyo107-852, JAPAN.  
<https://www.tsumura.co.jp/>